## **Exhibit D**

## Experiment 1 (killing a laboratory E. coli strain)

Donor 1: (control) S17-1 carrying self-transmissible plasmid without colE3 (plasmid is TetR, donor A below)

Donor 2: S17-1 carrying a non self-transmissible plasmid with colE3 (plasmid is TetR, donors B and C below)

Donor 3: S17-1 carrying a self-transmissible plasmid with colE3 (plasmid is T tR, donors D and E below)

Note: S17-1 is an E. coli strain carrying RK2 integrated in the genome

Recipient: CON4-54A (E. coli VH1000, RifR)

Donor and recipient cells were mixed and conjugated on a filter for 1 hour at 37°C, and exconjugants were selected by two antibiotics, Tet and Rif. Under this antibiotic selection, the donor or the recipient itself was not able to grow on this plate, which was shown at the bottom of each plate.

Conjugation plate (LB with Tet/Rif) selecting exconjugants




Table. Number of exconjugants counted from the plate above

dilution	Control (A)	non self-transmissible vector		self-transmissible vector	
		clone #1 (B)	clone #2 (C)	clone #1 ( <b>D</b> )	clone #2 ( <b>E</b> )
1	TNC	TNC	TNC	2	5
10 <sup>-2</sup>	TNC	19	12	1	1
10 <sup>-4</sup>	TNC	0	0	0	0
10 <sup>-6</sup>	5	0	0	0	0

NOTE: TNC - Too Numerous to Count

## Comparison of survival values after conjugation (based on the data shown above)

